

Defence Seminar

Seminar Title	: Role of functional amyloid and nucleotide second messenger in the marine bacterium <i>Pseudomonas aeruginosa</i> PFL-P1 for biodegradation of polycyclic aromatic hydrocarbon
Speaker	: Swetambari Kumari (Rollno : 518ls1002)
Supervisor	: Surajit Das
Venue	: Civil Engineering Department Seminar Room
Date and Time	: 09 Jun 2025 (11.00 AM)
Abstract	: The thesis illustrates the interaction of functional amyloid in <i>Pseudomonas</i> (Fap) with pyrene, a commonly encountered PAH. <i>Pseudomonas aeruginosa</i> PFL-P1, a marine biofilm-forming bacterium, showed strong tolerance (2200 ppm) and chemotaxis toward pyrene, indicating its potential for bioremediation in PAH-contaminated environments. The expression of <i>fapC</i> was upregulated six fold ($p < 0.0001$) when pyrene was used as the sole carbon source. Molecular docking simulations revealed a strong binding affinity between FapC and pyrene, with a binding energy of -6.75 kcal/mol, indicating a robust interaction. Confocal laser scanning microscopy (CLSM) further demonstrated a significant increase in amyloid production during biofilm formation by <i>P. aeruginosa</i> PFL-P1 in the presence of pyrene ($p < 0.0001$). The elevated expression of <i>fapC</i> and the hydrophobic interactions between FapC and PAH compounds indicate the critical role of Fap in PAH binding, facilitating their subsequent degradation. The study also explored the role of cyclic di-GMP (c-di-GMP), a bacterial nucleotide second messenger that plays a central role in regulating biofilm formation and stability. This study also reported the characterization of Fap fibrils from <i>P. aeruginosa</i> PFL-P1 and their interaction with pyrene, further assessing the impact on pyrene degradation. Overexpression of <i>fap</i> in <i>E. coli</i> BL21(DE3) cells enhanced biofilm formation ($p < 0.0001$) and amyloid production ($p = 0.0002$), particularly in the presence of pyrene. Defibrillated Fap analysis revealed FapC monomers, with increased fibrillation upon exposure to pyrene. Circular Dichroism (CD), Fourier Transform Infrared Spectroscopy (FTIR), and X-ray Diffraction (XRD) confirmed characteristic amyloid peaks and structural changes in Fap fibrils upon pyrene exposure. Three-dimensional Excitation-Emission Matrix (3D-EEM) analysis identified a protein-like fluorophore in Fap fibrils that exhibited fluorescence quenching upon pyrene binding. The binding constants ranged from 5.23 to 7.78 M^{-1} , with a ΔG of -5.10 kJ/mol at 298K, indicating a spontaneous, endothermic interaction primarily driven by hydrophobic forces. Exogenous Fap fibrils significantly enhanced biofilm growth and pyrene degradation by <i>P. aeruginosa</i> PFL-P1, increasing the degradation rate from 46% to 64% within seven days ($p = 0.0236$). Gas Chromatography Mass Spectrometry (GC-MS) analysis identified various metabolites, suggesting the involvement of the phthalic acid pathway in pyrene degradation. This study provides a deeper understanding of the structural dynamics of Fap fibrils in response to pyrene exposure and offers potential applications in environmental bioremediation, particularly for the degradation of PAHs.